

Upper Bound on the Hadronic Light-by-Light Contribution to the Muon $g - 2$

Jens Erler and Genaro Toledo Sánchez

Instituto de Física, Universidad Nacional Autónoma de México, 01000 México D.F., México

(Dated: February 2, 2008)

There are indications that hadronic loops in some electroweak observables are almost saturated by parton level effects. Taking this as the hypothesis for this work, we propose a genuine parton level estimate of the hadronic light-by-light contribution to the anomalous magnetic moment of the muon, $a_\mu^{\text{LBL}}(\text{had})$. Our quark mass definitions and values are motivated in detail, and the simplicity of our approach allows for a transparent error estimate. For infinitely heavy quarks our treatment is exact, while for asymptotically small quark masses $a_\mu^{\text{LBL}}(\text{had})$ is overestimated. Interpolating, this suggests to quote an upper bound. We obtain $a_\mu^{\text{LBL}}(\text{had}) < 1.59 \times 10^{-9}$ (95% CL).

PACS numbers: 13.40.Em, 14.60.Ef, 12.20.Ds, 12.38.Bx.

The E-821 Collaboration at BNL [1] has measured the anomalous magnetic moment of the muon, $a_\mu \equiv (g_\mu - 2)/2$, with an uncertainty of $\Delta a_\mu = \pm 0.63 \times 10^{-9}$. This would provide sensitivity to new physics scales, Λ , as large as $\Lambda \sim m_\mu / \sqrt{\Delta a_\mu} = 4.2$ TeV, where m_μ is the mass of the muon [2]. A new experiment [3] at BNL is approved aiming at even greater (± 0.2 ppm) precision and implying a reach up to 7 TeV. Unfortunately, the interpretation of a_μ is compromised by large theoretical uncertainties introduced by hadronic effects diluting the new physics sensitivity. These mainly arise from two-loop vacuum polarization (VP) effects, $a_\mu^{\text{VP}}(2, \text{had})$, and from the three-loop contribution of light-by-light type, $a_\mu^{\text{LBL}}(\text{had})$. The calculations of $a_\mu^{\text{LBL}}(\text{had})$ based on chiral perturbation theory [4, 5] are the only ones to date solidly based on a systematic expansion. However, the estimated uncertainty, $\gtrsim \pm 10^{-9}$, is significantly larger [5] than the measurement error. There is a variety of model estimates, all supplementing the dominant π^0 -exchange contribution with other resonance exchanges and $\pi^\pm$ -loops. Current analyzes [6, 7] agree reasonably well on the magnitude (where residual differences are largely understood) and the sign of $a_\mu^{\text{LBL}}(\text{had})$. They have reached a high level of sophistication, but the error estimates remain rough guesses. There is also an estimate [8] based on the instanton liquid model with a small ($< 10\%$) quoted error.

In this work we offer a complementary way to estimate $a_\mu^{\text{LBL}}(\text{had})$, with no need to commit to the dominance of any particular type of contribution. It is a naïve parton level estimate which is solid in the heavy quark limit where it matches onto perturbative QCD, but overestimates the light-by-light contribution in the chiral limit. Therefore, our approach naturally implies an *upper bound* for $a_\mu^{\text{LBL}}(\text{had})$, which is very useful in view of the a_μ measurement lying above the Standard Model prediction. It can also be applied to $a_\mu^{\text{VP}}(\text{had})$, which serves as a reference case and allows for a *transparent* error estimate.

Attempting to obtain $a_\mu^{\text{LBL}}(\text{had})$ directly at the parton level seems hopeless at first since perturbative QCD cannot be applied except for the heavy c and b quarks.

Moreover, in the chiral limit, in which the bulk of the effect arises from light, charged, and essentially point-like pseudo-Goldstone bosons, one has $a_\mu^{\text{LBL}}(\text{had}) < 0$ from scalar QED [9], while leptons and heavy quarks contribute positively [10]. However, as shown in Ref. [7], the $\pi^\pm$ -loops receive large chiral symmetry breaking corrections, rendering the net $\pi^\pm$ -contribution almost negligible. Thus, at least as far as $a_\mu^{\text{LBL}}(\text{had})$ is concerned, we are dealing with a kinematic regime that conceivably admits an alternative description in terms of quark in place of chiral degrees of freedom. Indeed, an estimate [7] of the typical momenta, μ , in the $\pi^\pm$ -loops yields $\mu \sim 4.25 m_\pi \approx 0.6$ GeV, *i.e.* large enough to feel individual partons even within pions. These observations revive the question whether hadronic light-by-light diagrams are best described by hadrons or by partons or a mix. We are aware of three parton based attempts in the literature. An early approach [11] using exclusively constituent quarks [12] resulted in a numerical estimate, albeit with the wrong sign. Ref. [13] opted for a mixed treatment in which constituent quark masses, $m_{u,d} = 0.33$ GeV, $m_s = 0.5$ GeV, and $m_c = 1.87$ GeV (with assumed errors of $\pm 10\%$), were employed in the analytical result of Ref. [10] and produced the correct sign. The adopted definition, values, and uncertainty were not further justified, but in any case the constituent quarks were assumed to yield only the high momentum contribution which was supplemented with π^0 -exchange graphs as described by the chiral constituent quark model [14]. Ref. [15] is very close to both our strategy and philosophy and will be commented on below relation (10).

Indeed, a main obstacle for any model involving quark degrees of freedom at hadronic scales is to find the appropriate quark mass *definition*. In contrast to pole masses, $\overline{\text{MS}}$ masses, $\hat{m}_q(\mu)$ (where μ is the renormalization scale), tend to produce small perturbation coefficients (at least for sufficiently inclusive quantities). We will therefore avoid pole masses for heavy quarks and similarly constituent masses for light quarks. The scale choice is implied by the requirement of avoiding spurious logarithms, $\ln \mu^2 / \hat{m}_q^2(\mu)$ or $\ln \mu^2 / m_\mu^2$. They can be nullified

(resummed) by choosing $\mu = \hat{m}_q(\mu)$ for $\hat{m}_q(\hat{m}_q) > m_\mu$ and $\mu = m_\mu$ for $\hat{m}_q(\hat{m}_q) < m_\mu$, *i.e.* the higher of the two masses sets the scale. There are also genuine logarithms of the form $\ln \hat{m}_q^2(\mu)/m_\mu^2$ which cannot be eliminated.

But for the light (u, d, s) quark contribution to $g_\mu - 2$, the values of $\hat{m}_q(\hat{m}_q)$ or $\hat{m}_q(m_\mu)$ are unknown because $\hat{\alpha}_s$ and consequently the anomalous quark dimensions formally diverge. In a similar context, however, it was possible to proceed phenomenologically. Namely, the hadronic contribution to the renormalization group evolution (RGE) of the QED coupling (the running), $\hat{\alpha}(\mu)$, can be mimicked [16] by effective quark masses. These have been defined more precisely in Ref. [17] as *threshold quark masses*, $\bar{m}_q$, and correspond to the scale at which the theory is changed to include or exclude the quark q and where the one-loop QCD β function coefficient changes correspondingly. Since this is *not* a perturbative description, higher order QCD effects are assumed to be absorbed in the threshold masses, leading to [17],

$$\bar{m}_u = \bar{m}_d = 176 \pm 9 \text{ MeV}, \quad (1)$$

$$\bar{m}_s = 305_{-65}^{+82} \text{ MeV}, \quad (2)$$

$$\bar{m}_c = 1.18 \pm 0.04 \text{ GeV}, \quad (3)$$

$$\bar{m}_b = 4.00 \pm 0.03 \text{ GeV}, \quad (4)$$

where isospin symmetry, $\hat{m}_u(\hat{m}_u) = \hat{m}_d(\hat{m}_d)$, was imposed. The uncertainties in the first (last) two lines have a correlation of -100% ($+29\%$). The values (1–4) are implied by the considerations [17] in the next paragraph.

Heavy quark masses are qualitatively given by one half of the mass, M_{1S} , of the corresponding $1S$ quarkonium resonance. If the quark is heavy enough that QCD perturbation theory can be applied, this statement can be made quantitative resulting in well studied QCD sum rules [18, 19], which are dominated by the $1S$ -quarkonium resonance and supplemented by higher vector resonances and a continuum contribution. They yield expressions for $\hat{m}_q(\hat{m}_q)$ with small QCD corrections which are known [20] and included up to order α_s^2 . The threshold masses in Eqs. (3–4) can then be obtained in terms of these using Eq. (32) in Ref. [17]. Defining the parameter, $\xi_q = 2\bar{m}_q/M_{1S}$ [17], we have that asymptotically $\xi_q \rightarrow 1$ for $\bar{m}_q \rightarrow \infty$ and $\xi_q \rightarrow 0$ for $\bar{m}_q \rightarrow 0$. Thus, for a heavy quark $\xi_q \sim 1$, while for a light quark $\xi_q \ll 1$. Also, we expect $\bar{m}_1 < \bar{m}_2 \implies \xi_1 < \xi_2$, which can be checked explicitly by comparing $\xi_b = 0.845 \pm 0.006$ with $\xi_c = 0.76 \pm 0.03$. This implies the upper bound $\xi_s < \xi_c$. A lower bound, $\xi_s > 0.47$, can be obtained by considering the $SU(3)$ limit, $\xi_u = \xi_d = \xi_s$. Since the total contribution to the running of α due to u, d , and s quarks is known experimentally [34], this (together with isospin symmetry) implies Eq. (2). Assuming in addition the suppression of flavor singlet contributions (the OZI rule [35]) then implies Eq. (1) and explains the complete anti-correlation with Eq. (2). Isospin violations will, of course, induce a small mass difference, $\bar{m}_d - \bar{m}_u$.

In Ref. [17] the threshold masses (1–4) were used for the RGE of the weak mixing angle, $\sin^2 \hat{\theta}(\mu)$, which is rigorously justified as the same vector current correlator convoluted by the same weight function enters in both, $\hat{\alpha}(\mu)$ and $\sin^2 \hat{\theta}(\mu)$. What we are proposing here, is that $a_\mu^{\text{LBL}}(\text{had})$ can be *modeled* by treating quarks like massive leptons and using the values (1–4). Notice, that these values are systematically lower than typical constituent masses [12], which is reasonable since we do not add an independent contribution from pion degrees of freedom.

It is *a priori* not obvious that this procedure and these values for the quark masses (in fact, any set of values) will generate the correct $a_\mu^{\text{LBL}}(\text{had})$, but this represents the only *ad hoc* model assumption that we need to make. One justification is the observation mentioned before, that the relevant kinematic region conceivably admits a description in terms of quark degrees of freedom, which is reinforced by the lower bounds, $\xi_s > 0.47$ and $\xi_{u,d} > 0.42$, showing that the physical light quark masses are midway between the perturbative ($\xi \rightarrow 1$) and chiral ($\xi \rightarrow 0$) limits. Thus, at a *qualitative* level, both $a_\mu^{\text{LBL}}(\text{had})$ and the RGE of $\hat{\alpha}(\mu)$ appear to be describable by moderately heavy quark degrees of freedom.

Although our treatment clearly improves with growing $\bar{m}_q$ and becomes exact for $\bar{m}_q \rightarrow \infty$, it is important to establish *quantitative* tests and to apply our approach to physical quantities for which the answer is known with sufficient precision. For this we note that $a_\mu^{\text{VP}}(2, \text{had})$ is described by the same vector current correlator as the RGE of $\hat{\alpha}(\mu)$ but is convoluted with a different weight function, $K^{(1)}(s)$ [21], so that just as in the case of $a_\mu^{\text{LBL}}(\text{had})$ it is *a priori* not obvious that our approach and quark mass values will generate the correct answer,

$$a_\mu^{\text{VP}}(2, \text{had}) \times 10^9 = 70.30 \pm 0.43 \pm 0.88 = 70.3 \pm 1.0 \quad (5)$$

To arrive at Eq. (5) we averaged the results based on e^+e^- annihilation data and τ decay data which are both taken from Ref. [22]. The second uncertainty accounts for the discrepancy between the two data sets. It is chosen to one half of the difference in central values and added in quadrature to the first (experimental) one. We compare this result with the analytical expression [23, 24],

$$a_\mu^{\text{VP}}(2, \text{had}) = \frac{\alpha^2}{\pi^2} N_C \sum_q Q_q^2 \sum_{n=0}^{\infty} \left(\frac{m_\mu}{\bar{m}_q} \right)^{2n+2} \times \quad (6)$$

$$\left[\frac{n \ln(m_\mu^2/\bar{m}_q^2)}{(n+3)(2n+3)(2n+5)} - \frac{8n^3 + 28n^2 - 45}{[(n+3)(2n+3)(2n+5)]^2} \right],$$

yielding,

$$a_\mu^{\text{VP}}(2, \text{had}) = 65.1_{-4.2}^{+2.6} \mp 0.1 \times 10^{-9}, \quad (7)$$

The first and second uncertainty are induced, respectively, from Eqs. (1–2) and Eqs. (3–4). Effects due to

$\bar{m}_d \neq \bar{m}_u$ cancel to first order in $a_\mu^{\text{VP}}(2, \text{had})$ (but not in $a_\mu^{\text{LBL}}(\text{had})$ below) and can be neglected. Notice, that the estimate (7) is in reasonable agreement with Eq. (5) and that our central value is *below* it. This is consistent with the chiral limit, $m_\pi \rightarrow 0$ with m_μ/m_π fixed, underestimating [36] $a_\mu^{\text{VP}}(2, \text{had})$. Indeed, in this limit,

$$m_\pi \rightarrow 0 \implies \frac{\bar{m}_q}{\mu} \rightarrow \left(\frac{m_\pi}{\mu}\right) \sqrt{\frac{1}{4N_C(Q_u^2 + Q_d^2)}} \approx \left(\frac{m_\pi}{\mu}\right)^{0.39},$$

(where μ is a fixed reference scale) so that $m_\mu/\bar{m}_q \rightarrow 0$. Combined with $K(s) > 0$ this implies a systematic *underestimate*. The fact that the estimate (7) reproduces the experimental result within about 10% provides phenomenological support for our approach and its use of quark degrees of freedom. We can also directly compare the contribution from c quarks, $a_\mu^{\text{VP}}(2, c) = 1.27 \times 10^{-9}$ from Eqs. (3) and (6), with the analytical result up to order α_s^2 originally obtained in the second Ref. [20] in the form given in Eq. (4) of Ref. [25]. Using $\hat{m}_c(\hat{m}_c) = 1.29$ GeV and $\hat{\alpha}_s(M_Z) = 0.1216 \pm 0.0017$ (corresponding to $\hat{\alpha}_s(\hat{m}_c) = 0.432$ and $\hat{\alpha}_s(\bar{m}_c) = 0.473$), the latter yields $a_\mu^{\text{VP}}(2, c) = 1.46 \times 10^{-9}$. Thus, here we underestimate the known result by 13%, which is traceable to different α_s dependences in the RGE of $\hat{\alpha}(\mu)$ and the leading $\mathcal{O}(m_\mu^2/\hat{m}_c^2)$ term in $a_\mu^{\text{VP}}(2, \text{had})$. QCD corrections to the $\mathcal{O}(m_\mu^4/\hat{m}_q^4)$ term which are significant only for lighter quarks tend to have a compensating effect, offering a perturbative rationale why our approach seems to work better for light quarks than for charm.

The success of the test in the previous paragraph may be coincidental. Therefore, we performed three similar tests on various three-loop VP effects, $a_\mu^{\text{VP}}(3i, \text{had})$, each with a different kernel function, $K^{(2i)}(s)$ [26, 27]. $i = a$ contains an additional photonic correction or muon-loop relative to $a_\mu^{\text{VP}}(2, \text{had})$, while $i = b$ (c) corresponds to an additional electron (hadron) loop insertion. Our method (in parentheses) reproduces the central values of Ref. [28],

$$\begin{aligned} a_\mu^{\text{VP}}(3a, \text{had}) &= -2.11 \times 10^{-9} \quad (-1.82 \times 10^{-9}), \\ a_\mu^{\text{VP}}(3b, \text{had}) &= +1.07 \times 10^{-9} \quad (+0.99 \times 10^{-9}), \\ a_\mu^{\text{VP}}(3c, \text{had}) &= +2.7 \times 10^{-11} \quad (+2.8 \times 10^{-11}), \end{aligned} \quad (8)$$

within 14%, 7%, and 4%, respectively. Note, that these represent genuine “hit or miss” tests involving no adjustable parameters, and all four passed at the 10% level.

We finally apply our approach to $a_\mu^{\text{LBL}}(\text{had})$. Our master formula is the one for heavy leptons [10],

$$a_\mu^{\text{LBL}}(\text{had}) = \frac{\alpha^3}{\pi^3} N_C \sum_q Q_q^4 \sum_{n=1}^{\infty} \frac{m_\mu^{2n}}{\bar{m}_q^{2n}} \sum_{k=0}^2 C_{n,k} \ln^k \frac{m_\mu}{\bar{m}_q^2},$$

where $C_{1,0} = 3/2\zeta_3 - 19/16$, $C_{1,1} = C_{1,2} = 0$, and where the other $C_{n,k}$ up to $n = 5$ can be found in Eq. (5) of Ref. [9]. The central values in Eqs. (1–4) then imply $a_\mu^{\text{LBL}}(\text{had}) = 1.36 \pm 0.13 \times 10^{-9}$, where the uncertainty

is the model error, determined by the 9.5% difference between the upper end of the range (5) from the central value in Eq. (7). Similarly, the upper and lower error in Eqs. (1–2) and (7) correspond to $a_\mu^{\text{LBL}}(\text{had}) = 1.27 \pm 0.18 \times 10^{-9}$ and $a_\mu^{\text{LBL}}(\text{had}) = 1.47 \pm 0.04 \times 10^{-9}$, *i.e.* model errors of 14.1% and 2.9%, respectively. As expected, b quark effects are negligible, but the contribution from c quarks (usually ignored) turns out to be 0.04×10^{-9} . Although small it might serve as an interesting testing tool of other model calculations of $a_\mu^{\text{LBL}}(\text{had})$ which should be consistent with what we have obtained, since our method is solid (up to effects of $\mathcal{O}(\alpha_s)$) for heavy quarks. To account for isospin breaking we shift $\bar{m}_d$ by +2 MeV and $\bar{m}_u$ by –1 MeV, increasing all results by 0.01×10^{-9} to

$$a_\mu^{\text{LBL}}(\text{had}) = 1.37_{+0.15}^{-0.27} \times 10^{-9}, \quad (9)$$

where the error covers the three ranges above and is constructed to include the intrinsic model uncertainty. As in any hadronic model there would be additional model uncertainty if the comparison to other cases (here $a_\mu^{\text{VP}}(\text{had})$) was regarded as pure coincidence. The result, $1.36 \pm 0.25 \times 10^{-9}$, of Ref. [7] is the first to take short-distance QCD constraints explicitly into account. By construction, this is a feature of our approach, as well, so it is gratifying that the estimate (9) turns out to agree perfectly. Our central value is *higher* than previous ones [6], which is expected given that in the chiral limit, $m_\pi \rightarrow 0$ with m_μ/m_π fixed (and presumably also with m_μ fixed, but the scalar QED result in this case is not known), we *overestimate* $a_\mu^{\text{LBL}}(\text{had})$ since $\pi^\pm$ -loops contribute negatively.

Now we repeat the above analysis with the model errors multiplied by 1.645 to shift from uncertainties estimated in a “1 σ spirit” to 90% ranges, and find $a_\mu^{\text{LBL}}(\text{had}) = 1.37 \pm 0.21 \times 10^{-9}$, $a_\mu^{\text{LBL}}(\text{had}) = 1.28 \pm 0.30 \times 10^{-9}$, and $a_\mu^{\text{LBL}}(\text{had}) = 1.48 \pm 0.07 \times 10^{-9}$, respectively. The upper errors have converged, and since as mentioned our approach tends to overestimate $a_\mu^{\text{LBL}}(\text{had})$, we quote as our final result the 95% CL upper bound,

$$a_\mu^{\text{LBL}}(\text{had}) < 1.59 \times 10^{-9}. \quad (10)$$

In a similar paper [15], a 1-parameter fit to $a_\mu^{\text{VP}}(2, \text{had})$ resulted in $m_u = m_d = m_s - 180$ MeV = 166 ± 1 MeV. These values are within errors consistent with Eqs. (1–2), reflecting our statement that $a_\mu^{\text{VP}}(2, \text{had})$ and $\hat{\alpha}(\mu)$ can be approximated using the same masses. We benefited from a deeper m_μ^2/m_q^2 -expansion of a_μ^{LBL} [9], lowering the light quark contribution relative to [15] by $\sim 6\%$. The most important new aspect here is that the recent scalar QED calculation [9] allows us to interpolate between the chiral and heavy quark limits. If one can prove (conceivably by lattice simulations) the monotony of the product $m_q^2 a_\mu^{\text{LBL}}(\text{had})$ as a function of m_q , our bound (10) would become rigorous, but this remains a loophole for now.

We stress that it is worthwhile to compute the α_s correction to the heavy quark light-by-light contribution to a_μ to check explicitly its magnitude and more importantly its sign, to compare it to the RGE of $\hat{\alpha}(\mu)$, and to use it as a refinement of our method. This contribution may be called $A_2^{(6,2)}(m_\mu/\hat{m}_q)$ and corresponds to a subclass of 60 of the 469 Feynman diagrams contributing at $\mathcal{O}(\alpha^4)$ to what is known as $A_2^{(8)}(m_\mu/m_\tau)$ with m_τ replaced by $\hat{m}_q(\hat{m}_q)$ and one prefactor of α by $\hat{\alpha}_s(\hat{m}_q)$. This is possible with state-of-the-art computer codes [29].

We have developed (quark mass definition, scale settings, quark mass values, and uncertainties) what we believe are necessary ingredients for a parton model estimate of $a_\mu^{\text{LBL}}(\text{had})$ and similar quantities. The model has four key benefits: (i) it is simple and transparent, (ii) it can be tested, as we did successfully using $a_\mu^{\text{VP}}(\text{had})$, (iii) short-distance constraints are taken into account by construction, and (iv) it naturally provides an upper bound on $a_\mu^{\text{LBL}}(\text{had})$ because, as we have argued, our method tends to yield an overestimate and also because the 95% upper error is virtually insensitive to our input masses.

We would like to compare our approach with celebrated models of low-energy QCD, such as the vector meson dominance model [30], the $1/N_C$ expansion [31], the chiral constituent quark model [14], or the extended [32] Nambu-Jona-Lasinio [33] model. It is not obvious *a priori* that approximations like $N_C = 3 \gg 1$ [31] or $\Lambda_{\text{QCD}} \ll M_{\chi_{SB}}$ (the spontaneous chiral symmetry breaking scale) [14], or the dominance of certain resonances [30] or operators [32] — while plausible — would generate useful results. But experience showed that they do. The assumption made in this work is that for the type of observables considered, parton level effects dominate and almost saturate hadronic loops. It is in line with the observations of Ref. [7], and we found further evidence for it in both, $a_\mu^{\text{VP}}(\text{had})$ and $a_\mu^{\text{LBL}}(\text{had})$, which passed quantitative tests with better than expected phenomenological success. This may be interpreted as pure coincidence and, clearly, it is important to submit our method to other case studies in the future. More optimistically, however, our results offer an alternative view on hadronic loops and may lead to a rigorous bound on $a_\mu^{\text{LBL}}(\text{had})$.

It is a pleasure to thank Michael Ramsey-Musolf and Lee Roberts for helpful comments on the manuscript and Lorenzo Díaz-Cruz and Peter Zerwas for fruitful discussions. JE greatly acknowledges the hospitality of Caltech, DESY (Hamburg), and U. Penn. This work was supported by CONACyT (México) contract 42026-F and by DGAPA-UNAM contract PAPIIT IN112902.

[1] BNL-E-821 Collaboration: G.W. Bennett *et al.*, *Phys. Rev. Lett.* **92**, 161802 (2004).
 [2] For a review, see A. Czarnecki and W.J. Marciano, *Phys.*

Rev. **D64**, 013014 (2001).
 [3] BNL-E-821 and E-969 Collaborations: D.W. Hertzog *et al.*, *Nucl. Phys. Proc. Suppl.* **144C**, 191 (2005).
 [4] M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, *Phys. Rev. Lett.* **88**, 071802 (2002).
 [5] M. Ramsey-Musolf and M.B. Wise, *Phys. Rev. Lett.* **89**, 041601 (2002).
 [6] M. Knecht and A. Nyffeler, *Phys. Rev.* **D65**, 073034 (2002); J. Bijnens, E. Pallante and J. Prades, *Nucl. Phys.* **B626**, 410 (2002); M. Hayakawa and T. Kinoshita, *Phys. Rev.* **D57**, 465 (1998).
 [7] K. Melnikov and A. Vainshtein, *Phys. Rev.* **D70**, 113006 (2004).
 [8] A.E. Dorokhov, *Acta Phys. Polon.* **B36**, 3751 (2005).
 [9] J.H. Kühn, A.I. Onishchenko, A.A. Pivovarov and O.L. Veretin, *Phys. Rev.* **D68**, 033018 (2003).
 [10] S. Laporta and E. Remiddi, *Phys. Lett.* **301B**, 440 (1993).
 [11] J. Calmet, S. Narison, M. Perrottet and E. de Rafael, *Rev. Mod. Phys.* **49**, 21 (1977).
 [12] A. De Rújula, H. Georgi and S.L. Glashow, *Phys. Rev.* **D12**, 147 (1975).
 [13] J.F. de Troconiz and F.J. Yndurain, *Phys. Rev.* **D65**, 093001 (2002).
 [14] A. Manohar and H. Georgi, *Nucl. Phys.* **B234**, 189 (1984).
 [15] A.A. Pivovarov, *Phys. Atom. Nucl.* **66**, 902 (2003).
 [16] W.J. Marciano and A. Sirlin, *Phys. Rev.* **D22**, 2695 (1980).
 [17] J. Erler and M.J. Ramsey-Musolf, *Phys. Rev.* **D72**, 073003 (2005).
 [18] V.A. Novikov *et al.*, *Phys. Rep.* **41**, 1 (1978).
 [19] J. Erler and M. Luo, *Phys. Lett.* **558B**, 125 (2003).
 [20] K.G. Chetyrkin, J.H. Kühn and M. Steinhauser, *Nucl. Phys.* **B482**, 213 (1996) and *ibid.* **505**, 40 (1997).
 [21] S.J. Brodsky and E. De Rafael, *Phys. Rev.* **168**, 1620 (1968).
 [22] M. Davier, A. Höcker and Z. Zhang, *hep-ph/0507078*.
 [23] H.H. Elend, *Phys. Lett.* **20**, 682 (1966).
 [24] M.A. Samuel and G. Li, *Phys. Rev.* **D44**, 3935 (1991).
 [25] J. Erler and M. Luo, *Phys. Rev. Lett.* **87**, 071804 (2001).
 [26] R. Barbieri and E. Remiddi, *Nucl. Phys.* **B90**, 233 (1975).
 [27] J. Calmet, S. Narison, M. Perrottet and E. de Rafael, *Phys. Lett.* **61B**, 283 (1976).
 [28] B. Krause, *Phys. Lett.* **390B**, 392 (1997).
 [29] T. Kinoshita and M. Nio, *Phys. Rev.* **D70**, 113001 (2004).
 [30] Y. Nambu and J.J. Sakurai, *Phys. Rev. Lett.* 879(1962).
 [31] G. 't Hooft, *Nucl. Phys.* **B75**, 461 (1974).
 [32] J. Bijnens, C. Bruno and E. de Rafael, *Nucl. Phys.* **B390**, 501 (1993).
 [33] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961) and *ibid.* **124**, 246 (1961).
 [34] The corresponding experimental error induces an additional uncertainty into $\bar{m}_u$, $\bar{m}_d$, and $\bar{m}_s$. This is an order of magnitude below the one in Eqs. (1–2) and neglected.
 [35] This assumption is not necessary to derive the $SU(3)$ limit, because in this case the flavor singlet contribution vanishes by virtue of $Q_u + Q_d + Q_s = 0$. OZI rule violations could still affect the result (1) but on general grounds we expect them to be tiny [17].
 [36] The alternative chiral limit, $m_\pi \rightarrow 0$ with m_μ fixed, is reproduced correctly, because then the leading effects in both $a_\mu^{\text{VP}}(2, \text{had})$ and the RGE of $\hat{\alpha}(\mu)$ are logarithmic.